



Trip Report

**The three Colombian Andes: Central, Western and Eastern Cordillera plus Chocó
bioregion and Cauca valley river specialties.**

Nov 19 to Dec 2/2018

Edwin Acevedo

Tour leader Native Birding Colombia



Gray-breasted mountain Toucan

This classic tour visited the well known and very good access places on Andes to enjoy of its rich avifauna, We were moving on premontane, humid, dry and highland forest and Paramo ecosystem. The weather was excellent with sunny days; but most of the time we were under shadow, different from the

first day along Porce area and day five in Bolombolo area near Cauca river which were hot; some rains at nights in some places and one morning in Tatamá N P; but did not affected our goals, birds were very active and the birders were so focused and skilled; transfers on time; everything flowed as desired; which is always our aim. Predominant bird Families were (Thraupidae) 79 taxa, Hummingbirds (Trochilidae) and Tyrant -Flycatchers 49 taxa each one. It was very ample about bird groups diversity. We shared lovely memories on field and everyone was satisfied !

This bird list follows broadly the taxonomy of South American Classification Committee SACC (2018). Each specie has the locality were it was recorded more easy; some of the birds could be seen in different spots.

Recorded taxa

DUCKS & GEESE @Anatidae

- Ruddy Duck (On Sumapaz lagoon) ´
- Torrent Duck - (Otún)
- Andean Teal - (On Sumapaz lagoon) ´

GUANS, CHACHALACAS & CURASSOWS: Cracidae

- Sickle-winged Guan - Chamaepetes goudotii (La Romera R)
- Andean Guan - Penelope montagnii (Hacienda El Bosque)
- Cauca Guan - Penelope perspicax Endemic/ Endangered (Otún Q)
- Wattled Guan - Aburria aburri Near Threatened
- Colombian Chachalaca - Ortalis columbiana Endemic (Piha reserve)

NEW WORLD QUAILS: Odontophoridae

- Chestnut Wood-Quail - Odontophorus hyperythrus Endemic - Near Threatened (Not easy anywhere)

GREBES Podicipedidae

- Least Grebe - Tachybaptus dominicus (On Sumapaz lagoon) ´
- Pied-billed Grebe - Podilymbus podiceps " " "

PIGEONS & DOVES: Columbidae

- Rock Pigeon - (Around towns)
- Band-tailed Pigeon - (Yellow -eared parrot R)
- White-tipped Dove - (Porce area)
- Eared Dove - Zenaida auriculata " "
- Ruddy Ground Dove - Columbina talpacoti " "

CUCKOOS: Cuculidae

- Smooth-billed Ani - *t x{nuamMMb* (Porce area)
- Squirrel Cuckoo - *<MMMMMM* (La Romera R)

NIGHTHAWKS & NIGHTJARS: Caprimulgidae

- Band-winged Nightjar - *@y{Zij} xMnl` bnyxy* (Río Blanco reserve)

SWIFTS: Apodidae

- White-collared Swift - *@xZu{nuxnM Z%b Mly* (Chicaque R)

HUMMINGBIRDS: Trochilidae

- White-necked Jacobin - *'jnxy|` Mk ZjbfnxM* (Piha reserve)
- Tawny-bellied Hermit - *<aM{anx byytk Mnuanx|y* (Río Blanco reserve)
- Green-fronted Lancebill - *ž nxt_ZxM| XnfbM* (One ind. in Piha reserve)
- Wedge-billed Hummingbird - *@aby{Zy` Zn_xntb* (One ind. Río Blanco reserve)
- Brown Violetear - *t njlxXZjuab M* (Río Blanco reserve)
- Lesser Violetear - *t njlx{aMMyb|y* « « «
- Sparkling Violetear - *t njlx{bvx| yMMy* (Chicaque R)
- Black-throated Mango - *° l {axMh{anxM.l b xhjjy* (Porce area)
- Amethyst-throated Sunangel - *) ZjbM` Zj|yM Z{aty{bhjjy* (Sumapaz NP)
- Tourmaline Sunangel - *) ZjbM` Zj|yZ. nx|y* (Yellow-eared Parrot reserve)
- Green Thorntail - *ž byhy| xMhl fZxyb* (One ind. Tatamá NP)
- Speckled Hummingbird - *° XZjnk tMk ZjMn` Zl ty* (Río Blanco reserve)
- Long-tailed Sylph - *°` jMhVxM|yi b` b* " " "
- Violet-tailed Sylph - *°` jMhVxM|yVhZjZy|y* Near endemic (Tatamá NP)
- Purple-backed Thornbill - *?Mk uank bxl k bxnatl Va|k* (Nevados NP)
- Rainbow-bearded Thornbill - *t aMMy{b k MaZxMib* « «
- Buffy Helmetcrest - *°° 5..tun` nl` y| UZjtb* Endemic Endangered (One indiv. in Nevado NP)
- Green-bearded Helmetcrest - *5..tun` nl` | Zx b* Endemic (One ind. in Sumapaz NP)
- Tyrian Metaltail - *2 Z{M| xM{tM{ab M* (Piha reserve)
- Greenish Puffleg - *) MjnuaMZXbM|xZjbM* (La Romera reserve)
- Black-thighed Puffleg - *" xM Zk byXZUt* Near endemic/threatened (Around hotel Termes)
- Golden-breasted Puffleg - *" xM Zk byk nyw ZxM* « «
- Shining Sunbeam - *°` jMNM{yV| uxwZl | by* « «
- Bronzy Inca - *t nZjb Zl MhZjb Zl M*
- Brown Inca - *t nZjb Zl M, bynl b* Near endemic (Easy in Tatama NP)
- Black Inca - *t nZjb Zl Mx| ZjZb* Endemic/Vulnerable/Near threatened (One ind. in Chicaque)
- Collared Inca - *t nZjb Zl M{nxw M* (Río Blanco reserve)
- Buff-winged Starfrontlet - *t nZjb Zl M| {Z{bM* (Around hotel Termes)
- Blue-throated Starfrontlet - *t nZjb Zl MaZjbM{aZM* Near endemic (Sumapaz NP)
- Mountain Velvetbreast - *1MZY MMJMZY Mb* (Chicaque R)

Sword-billed Hummingbird	- "I ybZxMVI ybZxM	(Hacienda El Bosque)
Great Sapphirewing	- <{ZxnuamIZyVtMnu{Zx}y	(Around hotel Termales)
Buff-tailed Coronet	- EnbynI ZMIMjMZYy	(very common and aggressive in Rio Blanco)
Velvet-purple Coronet	- EnbynI ZMIMMxb	Near endemic (Easy in Tatamá NP)
Booted Racket-tail	- 5VZM y I XZx, nnXb	(One ind. in Rio Blanco)
White-tailed Hillstar	- BxnVaxnMUn ` Zb	(Easy in Tatamá NP)
Purple-bibbed Whitetip	- Bxny{bZ UZI M b b	" "
Fawn-breasted Brilliant	-) ZjbnXn. Mx Ub nbZy	«- «-
Green-crowned Brilliant	-) ZjbnXn. MVM jM	(Piha reserve)
Empress Brilliant	-) ZjbnXn. Mx uZxMxb.	Near endemic (Easy in Tatamá NP)
White-bellied Woodstar	- t aM{nVZxM yk jyM{	(one ind. in Rio Blanco)
Gorgeted Woodstar	- t aM{nVZxM yaZjbnXn	(Jardín Encantado feeders)
Purple-throated Woodstar	- t Mjwajn. .k tVazjib	Near endemic (Easy in Tatamá NP)
Western Emerald	- t ajnxy{Uhl k ZjMnxtI Va y	«- «-
Crowned Woodnymph	- AaMj xMIMMhjnK UBM	(Tatamá NP)
Rufous-tailed Hummingbird	- ° k MjM{MVMj	(Porce area)
Andean Emerald	- ° k MjM{xMVMZ	(Piha reserve)
Steely-vented Hummingbird	- °° k MjM{MVMZxn{Zb	(Easy in Tatamá NP)
Indigo-capped Hummingbird	- ° k MjM{MVMb nl y	Endemic (Jardín Encantado feeders)

RAILS, CRAKES & COOTS: Rallidae

Bogota Rail	- ?Mj yyZk b j k UZ y	Endemic Endangered (Sumapaz NP)
Spot-flanked Gallinule	- <nxaatxuyk ZjMnu	(Sumapaz NP)
Common Gallinule	- ° (Mj b jM MZMM	«- «-
American Coot	- ° jMVMK ZxMIM	«- «-

PLOVERS: Charadriidae

Southern Lapwing	- GMIZj yVabZI yb	(Porce area)
------------------	-------------------	--------------

SANDPIPERS, SNIPES & PHALAROPES: Scolopacidae

Pectoral Sandpiper	- t Mj b yk ZjMn{ny	(Porce area)
Noble Snipe	- (Mj b MnI nUby	Near-threatened
Solitary Sandpiper	- Axo` Mj b kMVM	(Porce area)

ANHINGA & CORMORANTS: Phalacoracidae

Neotropic Cormorant	- Phalacrocorax brasilianus	(Porce area)
---------------------	-----------------------------	--------------

HERONS, EGRETS & BITTERS: Ardeidae

Black-crowned Night-Heron	- 3tV{b xM.I tV{b xM.	(Porce area)
Striated Heron	- t {nxZy{MVM	«- «-
Cattle Egret	- t U jV yby	

- Cocoi Heron - ° xXZMhVhb «← «←
- Great Egret - °° xXZMYUM
- Snowy Egret - °° xZ{{M[a]jM «← «←
- Little Blue Heron - °° xZ{{MMZx}jZM «← «←

IBISES AND SPOONBILLS: Threskiornithidae

- Bare-faced Ibis - <atk ny|y|b _|yMM|y (Porce area)

VULTURES AND CONDORS: Cathartidae

- Turkey Vulture - † MaMxZyMxM (Porce area)
- Lesser Yellow-headed Vulture - † MaMxZyU| xMfM|y «← «←
- Black Vulture - † nxM|tuyMxM|y (everywhere)
- Andean Condor - G|j|{x`xtua|y Near-threatened (Flying over Nevados N P)

OSPREY: Pandionidae

- Osprey - <MXbm| aN|MM|y (Porce area)

KITES, HAWKS & EAGLES: Accipitridae

- White-tailed Kite - ° jM|y|Z|V|x|y (Porce area)
- Swallow-tailed Kite - ° jM|nbZy_nx_bM|y (Piha and Tatamá NP)
- Black Hawk-Eagle - @|bM|{y|{tM|l|y (Tatamá NP)
- Sharp-shinned Hawk - °° WubZxy{xM|y (Otún Q.)
- Savanna Hawk - †|{Zn`M|y|k ZxM|N|y (Flying over Porce Area)
- Roadside Hawk - ?|unx|byk M|bny{x|y (Porce area)
- White-tailed Hawk - (ZM|nM|{y|yM|bM|xM|y (Río Blanco reserve)
- Black-chested Buzzard- ° M|jZ`° (ZM|nM|{y|k ZjM|njZ|V|y (Flying over Nevados NP)
- Gray-lined Hawk - †|{Zn|b|x|y (Porce Area)
- Broad-winged Hawk - †|{Zn|ujM|tu{Z|x|y (Yellow-eared Parrot reserve)

TYPICAL OWLS: Strigidae

- Tropical Screech - 5,, j`2 Z` M|VhuyVanj|UM (Piha R)
- Colombian Screech-5,, j`°°2 Z` M|VhuyVhjk UM|y (Tatamá NP)
- Andean Pygmy-Owl - (jM|Vx|k hM|x|b|b (Chicaque R)
- Stygian Owl - °° ym|y|t`b|y (Río Blanco reserve)

QUETZALS & TROGONS: Trogonidae

- Golden-headed Quetzal - <aMnk M|x|yM|x|Zuy (Río Blanco reserve)
- Collared Trogon - An`nl`VhjjM|y «← «←
- Masked Trogon - An`nl`uZxnl M|y (Tatamá NP)

MOTMOTS: Momotidae

- Tody Motmot -) tjnk M|Zy|k nk n{|jM (Porce Area)

Broad-billed Motmot - "jZV{xnl ujMtxatl Va|k (Porce Area)

Andean Motmot - 2 nk n{ | y iVzv{ Mnxvly (Río Blanco reserve)

KINGFISHERS: Alcedinidae

Ringed Kingfisher - 2 Z` NVZxtjZ {nxv{ MIM (Porce Area)

Green Kingfisher - t ajnxvZxtjZ MZ ZxvMIM «← «←

PUFFBIRDS: Bucconidae

Moustached Puffbird - 2 NjMhu{lpMk ty{Mvly (Cauca-Bolombolo area)

Lanceolated Monklet - 2 bxnk nl NvMjMvZnjMIM (Piha reserve)

BARBETS: Capitonidae

White-mantled Barbet - t Mtkn'atunjZ| V|y Endemic (Piha reserve)

Red-headed Barbet - "/ U| Wh Uh| xVZxb (Piha reserve)

TOUCAN BARBET: Semnornithidae

Toucan Barbet - @Zk l nx| byxNk ualMk|y Near Endemic (Tatamá NP)

TOUCANS: Ramphastidae

Emerald Toucanet - ° |jMhxatl Va|yuxVb|y (La Romera reserve)

Crimson-rumped Toucanet - ° |jMhxatl Va|yaMzk Minut` |y (Jardín area)

Gray-breasted Mountain-Toucan - ° l Xb Zl Matun` jMIM Near-threatened (Hacienda El Bosque)

Black-billed Mountain-Toucan - ° l Xb Zl M b xnyxy (Río Blanco reserve)

WOODPECKERS: Picidae

Olivaceous Piculet - <M|k l | ynjb{MZY|y (Porce area)

Grayish Piculet - <M|k l |y` xMIMZY|y Endemic (Montezuma Farm)

Acorn Woodpecker - 2 ZjMZXuZy_nxk b{fnx|y (Río Blanco)

Red-crowned Woodpecker - 2 ZjMZXuZy_x| UvMdtj|y (Porce area)

Smoky-brown Woodpecker - <VhbZY_|k b M|y (Tatamá NP)

Choco Woodpecker - GZl l{bnx| byVanVhZl yly Near Endemic/threatened (Tatamá NP)

Powerful Woodpecker - t Nk uZualj|yunjjZl y (Río Blanco reserve)

Lineated Woodpecker - ž xtnVhu|yjb ZM|y (Porce area)

Golden-olive Woodpecker - t njMk{Zy_x| Uv b ny|y (Jardín area)

Crimson-mantled Woodpecker - t njMk{Zy_xfjnb (Río Blanco reserve)

Spot-breasted Woodpecker - t njMk{Zy_u|l V{b|jM (Porce area)

FALCONS AND CARACARAS: Falconidae

Crested Caracara - t MvMvMvMvZxb M (Porce area)

Yellow-headed Caracara - 2 lɔʔMn Vatk Matak M «←» «←»
 American Kestrel - 'Nɪh yuMfZxɔy (Río Blanco reserve)

PARROTS: Psittacidae

Orange-chinned Parakeet - ɛxn{n Zxyh` | jMly (Porce area)
 Speckle-faced Parrot - <ɔnl | y'k | j'k | ny | y (Otún Q)
 Blue-headed Parrot - <ɔnl | yk Zl yx' | y (Porce area)
 Bronze-winged Parrot - <ɔnl | yValNhu{Zx' y «←» «←»
 Spectacled Parrotlet - ' nɪx | yVhl yubɔjM | y (Porce area)
 Brown-breasted Parakeet - <txa | xMMjɔkZxM Endemic Vulnerable/Endangered (Sumapaz)
 Brown-throated Parakeet - " | uy'k | jMuZx' b M (Porce area)
 Golden-plumed Parakeet - 1Zu{nɔk{NMMbM b M Vulnerable (Río Blanco reserve)
 Yellow-eared Parrot - 5` l nax' l Va | y b {Zn' by Endangered (its reserve)

ANTBIRDS: Thamnophilidae

Great Antshrike - ANMMk Mnx (Porce area)
 Bar-crested Antshrike - AaMk l nuab | yk | j'by{xM | y «←» «←»
 Black-crowned Antshrike - AaMk l nuab | yMx' | VaM (Cauca river Area)
 Uniform Antshrike - AaMk l nuab | y | l bhjnx (Piha reserve)
 Plain Antwreos - ž tykaMk l | yk Zl {Nly (Piha reserve)
 Bicolored Antwreos - ž tykaMk l | ynVxZl {Nly Near Endemic Vulnerable
 Checker-throated Antwren - " ub ZVnuatjM | j'fZl {xy (Porce area)
 Slaty Antwren - 2 tx n{ax' jMVab' bhjnx (Otún Q)
 Yellow-breasted - ° l {, xZl) Zuy' nVak | yMjMly Vulnerable
 Streak-headed Antbird - ž xtk nuab | y{xM bZuy (Río Blanco reserve)
 Parker's Antbird - † ZXhk MxnbZy uM' Zx' Endemic (Piha reserve)

ANTPITTAS & GNATEATERS: Conopophagidae, Grallariidea

Chestnut-crowned Gnateater - † nl nunuaMIMMx{M ZbZuy (Piha reserve)
 Chestnut-crowned Antpitta - (xjMxMk' bM bMjM (Río Blanco reserve)
 Bicolored Antpitta - (xjMxMk' nVb ZxM Near Endemic Vulnerable (Río Blanco)
 Rufous Antpitta - (xjMxMk' jM (Hacienda El Bosque)
 Tawny Antpitta - (xjMxMk' kZl y (Nevado NP)
 Brown-banded Antpitta - (xjMxMk' bZx Endemic Vulnerable Endangered
 Slate-crowned Antpitta - (xjMxMk' jM MIM (Nevado NP)
 Crescent-faced Antpitta - (xjMxMk' jM b Zbnl y Near Endemic Near-threatened (Hacienda El Bosque)

TAPACULOS: Rhinocryptidae

Ocellated Tapaculo - ° Vnu{Zx' bynx'anl t... (Tatamá NP)
 Blackish Tapaculo - @t{Nhu | yjMxMly (Río Blanco reserve)
 White-crowned Tapaculo - @t{Nhu | yMxM | y (Piha reserve)

Stiles's Tapaculo - @t{Nhu y bZyb	Endemic Endangered	(Piha reserve)
Nariño Tapaculo - @t{Nhu y b b bx	Near Endemic	(Tatamá NP)
Tatama tapaculo - @t{Nhu y NjMz%kauZ%b	Endemic SNE	" "
Pale-bellied Tapaculo - @t{Nhu y xyZbHjly	Near Endemic	Sumapaz NP

OVENBIRDS: Furnariidae

Tyrannine Woodcreeper - ž Zl XnVb VM{tMII b M		(Tatamá NP)
Strong-billed Woodcreeper - l wanVhjM{Zyuxnk ZnubatI Va y		(Otún Q)
Cocoa Woodcreeper - l wanxatl Va yy y xMIy	«	«
Olive-backed Woodcreeper - l wanxatl Va y{xMI jMly	«	«
Brown-billed Scythebill - Campylorhamphus pusillus		(Piha reserve)
Streak-headed Woodcreeper - l ZubXnVhjM{Zyyn jZtZ{b	«	«
Montane Woodcreeper - l ZubXnVhjM{ZyjMxtk b Zx		(Río Blanco reserve)
Streaked Xenops - l Zl nuyx {bMly	«	«
Buffy Tuftedcheek - <yZ XnVhjM{ZyjM xZl Vb		(Tatamá NP)
Streaked Tuftedcheek - <yZ XnVhjM{ZyUnbynl l ZM {b		(Río Blanco reserve)
Chestnut-winged Cinclodes - l b VjnXZy VUxbZl {xy		(Sumapaz NP)
Stout-billed Cinclodes - l b VjnXZyZ.VZjybx		(Nevados NP)
Slaty-winged Foliage-gleaner - <ab{Xnx_ yMzI l Z		(Porce area)
Buff-fronted Foliage-gleaner - <ab{Xnx_ k		(Piha reserve)
Montane Foliage-gleaner - ° l MUMZx{abM{xbMbhjly	«	«
Scaly-throated Foliage-gleaner - ° l MUMZx{abM{MZ MbZuy		(Tatamá NP)
Lineated Foliage-gleaner - @l XMM{tjM UNjMly		(Piha reserve)
Flammulated Treehunter - AaxwMZXV{Zy_jMk jM y		(Tatamá NP)
Striped Treehunter - AaxwMZXV{Zyanjny{b y		(Piha reserve)
Spotted Barbtail - Premnoplex brunnescens		(La Romera reserve)
Fulvous-dotted Treerunner - 2 M Mnx by{xZjM y	Near-threatened	(Tatamá NP)
Pearled Treerunner - 2 M Mnx by{w NR b Zx		(Río Blanco reserve)
Andean Tit-Spintail - l Zu{M{aZl xMVIxbHjM		(Nevados NP)
White-browed Spintail -) Zjk MxZM jMly		(Yellow-eared Parrot reserve)
Many-striped Canastero - ° y{aZl Zy_jMk jM y		(Sumapaz NP)
White-chinned Thistletail - ° y{aZl Zy_ jb b nyM		(Nevados NP)
Red-faced Spintail - l xM bjZ VZxt{axnuy		(Otun Q)
Silvery-throated Spintail - @l NjMly U XbM	Endemic	(Sumapaz NP)
Pale-breasted Spintail - @l NjMly VUZyZl y		(Porce area)
Azara's Spintail - @l NjMly VbM		(Río Blanco reserve)
Rufous Spintail - @l NjMly l bx _M		(Montezuma)

TYRANT - FLYCATCHERS: Tyrannidae

Black-capped Tyrannulet - <atjnk tby b xnVUbj y		(Río Blanco reserve)
Ashy-headed Tyrannulet - <atjnk tby Vb ZxZbZuy	«	«
Tawny-rumped Tyrannulet - <atjnk tby xnut bly		(Sumapaz NP)

Yellow-crowned Tyrannulet - 'AtxMI j yZjM y	(Porce area)
Yellow-bellied Elaenia - "jNI BIjMfn MZx	«← «←
Mountain Elaenia - "jNI BIxMI{?b	(Yellow-eared Parrot reserve)
White-tailed Tyrannulet - 2 ZhVZxM j yunZVnVZxM y	(Río Blanco reserve)
White-banded Tyrannulet - 2 ZhVZxM j yx(Bnu{Zx y	«← «←
White-throated Tyrannulet - 2 ZhVZxM j yjZ Vhuaxy	(Hacienda El Bosque)
Torrent Tyrannulet - Serpophaga cinerea	(Otún Q)
Bronze-olive Pygmy-Tyrant - Pseudotriccus pelzelni	(Tatamá NP)
Rufous-headed Pygmy-Tyrant - <yZ Xn{xM yx bZuy	(Rio Blanco reserve)
Golden-faced - Tyrannulet - Lt k Zx yVaxtynuy	(Porce area)
Marble-faced Bristle-Tyrant - <atjjnyM{Zynua{aYk B y	(La Romera Reserve)
Streak-necked Flycatcher - 2 bn ZVZy{xM bhjly	(Río Blanco reserve)
Ochre-bellied Flycatcher - 2 bn ZVZynjZM b Z y	(Porce area)
Rufous-breasted Flycatcher - 1Zu{nun' nl' x bZV y	(Otún Q)
Ornate Flycatcher - 2 tbn{xM yn M y	(Piha reserve)
Scale-crested Pygmy-Tyrant - 1nuan{xM yubZM y	(Piha reserve)
Black-throated Tody-Tyrant - ') Zk b{xM y' xMIMZI yb	(Yellow-eared Parrot reserve)
Rufous-crowned Tody-Flycatcher - <nZVn{xM yx bZuy	(Rio Blanco reserve)
Common Tody-Flycatcher - AnXbn{y x k Vb Zx k	(Porce area)
Fulvous-breasted Flatbill - ?atl VanVfV y jfawZV y	(Piha reserve)
Cinnamon Flycatcher - <txank tbnVb M nk Z y	?ah ÉjM Vh xZyZxZÁ
Handsome Flycatcher - '3ZuaZjnk tbn ul jVaZx	(Tatamá NP)
Acadian Flycatcher - "k ubXnl M. fozYVZ y	(La Romera Reserve)
Smoke-colored Pewee - t nl {nu y k b M y	(Tatamá NP)
Western Wood-Pewee - t nl {nu yynx x j y	(Tatamá NP)
Eastern Wood-Pewee - t nl {nu yf bZ y	(Tatamá NP)
Tropical Pewee - t nl {nu yVb ZxZ y	(La Romera Reserve)
Black Phoebe - @Mnx by b xM y	¿5{-I >Á
Vermilion Flycatcher - <txnVZuaM yx Ub y	(Porce area)
Streak-throated Bush-Tyrant - 2 tbn{aZxZ{Zy{xM bhjly	(Yellow-eared Parrot reserve)
Smoky Bush-Tyrant - 2 tbn{aZxZ{Zy k b M y	(Yellow-eared Parrot reserve)
Crowned Chat-Tyrant - 5Va{anZVMxnl {M y	(Río Blanco reserve)
Yellow-bellied Chat-Tyrant - 5Va{anZVMx k M	«← «←
Slaty-backed Chat-Tyrant - 5Va{anZVMb M nk ZbZ {xy	" "
Rufous-breasted Chat-Tyrant - 5Va{anZVMx bZV{nx y	(Nevados NP)
Brown-backed Chat-Tyrant - 5Va{anZVM k bhjnx	§ §
Cattle Tyrant - Machetornis rixosa	(Porce area)
Piratic Flycatcher - 'Z' M yjZ Vhuam y	«← «←
Rusty-margined - 'jtVMVaZx'2 tbn%Z{ZyVMIMZI yb	«← «←
Great Kiskadee - <éM y jua xM y	«← «←
Lemon-browed Flycatcher - t nl nub yVb Van Z{b	Vulnerable (Tatamá NP)
Golden-crowned Flycatcher - 2 tbnXtl M{ZyVaxtynVZuaM y	(Porce area)
Streaked Flycatcher - 2 tbnXtl M{Zyk M jM y	«← «←

- Tropical Kingbird - *AtxMI | y'k ZjM VanjM y* «[←]»[←]
 Fork-tailed Flycatcher - *AtxMI | y yM MIM* (Porce area)
 Pale-edged Flycatcher - *2 tM Ma | y VZuaNh{Zy* (Río Blanco reserve)

COTIGAS: Cotingidae

- Green-and-black Fruiteater - *<bxZnjMkZ_Zxb* (Río Blanco R)
 Barred Fruiteater - *<bxZnjM M MIM* (Hacienda el Bosque)
 Orange-breasted Fruiteater - *<bxZnjM h V | I XM* Near endemic
 Red-crested Cotinga - *° k uZjbnl xj UxnVxy{M | y* (Yellow-eared Parrot R)
 Andean Cock-of-the-rock - *? | ubhjMUZxj fM | y* (Jardín area)
 Purple-throated Fruitcrow - *> | Zxj jMu | xu | xMIM* (Porce area)
 Red-ruffed Fruitcrow - *Pyroderus scutatus* (Very common in Otún Q)
 Chestnut-capped Piha - *1wM | y, ZUZxb* Endemic Critically
 Dusky Piha - *1wM | y yM Vb ZxZ | y* (Río Blanco R)

MANAKINGS: Pipridae

- Yellow-headed Manakin - *t ajnxbun jM M M M M* (La Romera R)
 Golden-winged Manakin - *2 M h y Vaxtynu{Zx | y* (Tatamá NP)
 White-bearded Manakin - *2 MIM | yk MIM | y* (Porce Area)
 Club-winged Manakin - *2 M M Zxnu{Zx | y XZj M M y | y* (Tatamá NP)
 White-crowned Manakin - *ž bbat M M M* (Piha R)

TITYRAS, SCHIFFORNIS AND BECARDS: Tityridae

- Barred Becard - *<M atxMk ua | y fZxybhjnx* (Río Blanco R)
 Cinnamon Becard - *<M atxMk ua | y Vb I M nk Z | y* (Porce area)
 White-winged Becard - *<M atxMk ua | y unjt Vanu{Zx | y* (La Romera R)

VIREOS & GREENLETS: Vireonidae

- Black-billed Peppershrike - *t fV Maly | b xony{xy* (Otún Q)
 Rufous-naped Greenlet - *<M atytj fM Zk bx | I I ZM* «[←]»[←]
 Brown-capped Vireo - *GbxZnjZ | Vhuaxty* (Río Blanco)
 Red-eyed Vireo - *GbxZnjb M Z | y* (Otún Q)

JAYS: Corvidae

- Black-collared Jay - *t tM njt M M M k M M* Near endemic (Río Blanco)
 Beautiful Jay - *t tM njt M M | j VaxM* Near endemic, Threatened, Vulnerable (Tatamá NP)
 Black-chested Jay - *t tM nVhxM. M b by* (Piha reserve)
 Green Jay - *t tM nVhxM. t | Vxy* «[←]»[←]

SWALLOWS: Hirundinidae

- Blue-and-white Swallow - *< t nVáZjbnl VtM njZ | M* (Porce area)

Brown-bellied Swallow - 5xnVáZjbxnl k | xó M (Río Blanco)
 Southern Rough-winged Swallow - @Zj` bxnu{Zxt...x|_bñjby (Porce area)

WRENS: Troglodytidae

House Wren - Axn` jnXt{ZyVZXnl (Porce area)
 Mountain Wren - Axn` jnXt{Zyynjy{bñjby (Río Blanco)
 Sedge Wren - t by{n{anx| yujMZI yb (Nevados NP)
 Apolinar's Wren - t by{n{anx| yMñjtb Mb Endemic Endangered
 Sooty-headed Wren - <aZ|` nuZXb yyuVXb.. Near endemic
 Antioquia Wren - Aaxtnuab| yjZx Mb Endemic SNE
 Rufous Wren - t b| tVZx{abM| b|_M ¿AMM O3<Á
 Sharpe's Wren - t b| tVZx{abñjtbMZI y «- «-
 Gray-breasted Wood-Wren -) ZI bñxab MJ| Vhuaxy (La Romera)
 Munchique Wood-Wren -) ZI bñxab M Z xZ{b Endemic Critically endang/Vulnerable ¿AMM O3<Á

GNATCHATCHERS & GNATWRENS: Polioptilidae

Tropical Gnatcatcher - Polioptila plumbea (Bolombolo area)

DIPPER: Cinclidae

White-capped Dipper - t b V| yjZ| VñZuaM| y ¿5{-I >Á

DONACOBIOUS: Danocobiidae

Black-capped Donacobius - žnl MñUy yMñMñjM (Porce Area)

THRUSHES: Turdidae

Andean Solitaire - 2 tVZy{ZyXñjñbxZy (La Romera R)
 Slaty-backed Nightingale - Aax| ya t t MaM| y|_yMIZx «- «-
 Swainson's Thrush - t MaM| y| y{ jM| y (La Romera R)
 Black Solitaire - "I {nk nXZy{ZyVñXñb | y Near endemic (Tatamá NP)
 Pale-eyed Thrush - A| xX| yjZ| Vñuy (Otún Q)
 Black-billed Thrush - A| xX| yb| nUjy (Porce Area)
 Great Thrush - A| xX| y|_yMIZx (La Romera R)

MOCKINGBIRD: Mimidae

Tropical Mockingbird - 2 tk | y` bñf| y (Porce area)

TANAGERS: Thraupidae

White-capped Tanager - @ZñhyytuaMñUnWñM (Otún Q)
 Black-capped Hemispingus -) Zk bñb` | yMñmñZ| y (Río Blanco)
 Superciliaried Hemispingus -) Zk bñb` | y| uZññññy «- «-

Oleaginous Hemispingus -) Zk byub` y_xnl {Mly	«	«	
Black-eared Hemispingus -) Zk byub` yk ZjMn{ly	«	«	
Black-headed Hemispingus -) Zk byub` yfZx{Mly			
Gray-hooded Bush Tanager - t I Zk nyhu yx Uxny{xly	"	"	
White-lined Tanager - AMWtuan yx _l y			(Porce area)
Crimson-backed Tanager - ?Nk uanVZj yXtk bXbM y	"	"	
Flame-rumped Tanager - ?Nk uanVZj y_jNk k b Zx y			(Otún Q)
Black-and-gold Tanager - ŁMI` ybvk ZjMnVajNk ty	Endemic	Vulnerable	(Tatamá NP)
Gold-ringed Tanager - ŁMI` ybMlxZnVb VM	Endemic	Endangered	Vulnerable " "
Hooded Mountain-Tanager - Ł {axMlyk nl {MIM			(Hotel Termales area)
Black-chested Mountain-Tanager - t I Zk MaxMlybZ. tk bM			(Sumapaz NP)
Grass-green Tanager - t ajnrx byxZ_Zxb			(Yellow-eared Parrot R)
Lacrimose Mountain-Tanager - ° I byn` I Ma yjMxk ny y	«	«	«
Scarlet-bellied Mountain-Tanager - ° I byn` I Ma yb bZl {xly			(Hotel Termales area)
Blue-winged Mountain-Tanager - ° I byn` I Ma ynk u{ ny y			(Río Blanco)
Black-chinned Mountain-Tanager - ° I byn` I Ma y n{Mly	Near endemic		(Tatamá NP)
Buff-breasted Mountain-Tanager - ž U ybM{MIM			(Río Blanco)
Purplish-mantled Tanager - *Xnyx byunxuatxnVzuaM y	Near-threatened		(Tatamá NP)
Golden-crowned Tanager - *Xnyx byx _bZx{Z..			(Hotel Termales area)
Glistening-green Tanager - t ajnrxVaxtyMuanZl bh{ly	Near endemic		(Tatamá NP)
Multicolored Tanager - t ajnrxVaxtyM kXbytk M			Endemic Vulnerable
Blue-gray Tanager - AaxMlybZubhu y			
Palm Tanager - AaxMlyb y k M k			
Blue-capped Tanager - @unx{axMlybVtMnVzuaM			(Tatamá NP)
Golden-naped Tanager - AMI` MVMk bZyfb..			(La Romera R)
Black-capped Tanager - AMI` MVMaZb Zb	«	«	
Scrub Tanager - AMI` MVMf{xnb M	Near endemic		(Piha R)
Golden-hooded Tanager - AMI` MVMfMfM			(Porce area)
Blue-necked Tanager - AMI` MVMfMbhjly			(Otún Q)
Rufous-throated Tanager - *.n{axMlyb x _b jM	Near endemic		(Tatamá NP)
Speckled Tanager - *.n{axMlyb {MIM			(Piha R)
Blue-and-black Tanager - AMI` MVMfMynxb			?dh ŁjM Vh ?A
Beryl-spangled Tanager - AMI` MVM bxnfbxly	«	«	
Metallic-green Tanager - AMI` MVMfMfMnxXZy	«	«	
Plain-colored Tanager - AMI` MVMb nx MIM			(Porce area)
Bay-headed Tanager - AMI` MVM fxnjM			(Piha R)
Flame-faced Tanager AMI` MVMfMfMfM			(Tatamá NP)
Golden Tanager - AMI` MVMf{a y			(La Romera R)
Silver-throated Tanager - AMI` MVMf{ZnVzuaM			(Tatamá NP)
Swallow Tanager - AZyb Mfxbly			(Porce area)
Blue Dacnis - ž MI byMfMIM	«	«	
Purple Honeycreeper - t tMIZuZyVZx jZ y	«	«	
Green Honeycreeper - t ajnrxuaMIZyubM	«	«	

Guira Tanager -) Zk kaxMluby bM		(Jardín Area)
Scarlet-and-white Tanager - t axtn{ajtubyNyk nl b	Near endemic	(Piha R)
Blue-backed Conebill - t nl bny{x k yk{bhjnx		(Yellow-eared Parrot R)
Capped Conebill - t nl bny{x k NyUxnl y		(Río Blanco R)
Rufous-browed Conebill - t nl bny{x k x _ k	Near endemic	(Sumapaz NP)
Chestnut-bellied Flowerpiercer ž b jnyM jnxnybyk M	Endemic Endangered Vulnerable	(Tatamá NP)
Glossy Flowerpiercer - ž b jnyMjMZY Mtb		(Sumapaz NP)
Black Flowerpiercer- ž b jnyMa k ZxVly		(Chicaque R)
White-sided Flowerpiercer - ž b jnyMjUjMZXM		(Río Blanco R)
Indigo Flowerpiercer - ž b jnyMb Xb n{bM		(Piha R)
Rusty Flowerpiercer - ž b jnyMj{nXZy		(Río Blanco R)
Bluish Flowerpiercer - ž b jnyMMZx jZYZI y	« «	
Masked Flowerpiercer - ž b jnyMMtMIZM	« «	
Plushcap - t MjM UjxatI Va yXbXZk M		(Río Blanco R)
Plumbeous Sierra-Finch - <axt` b y I bhjnx		(Sumapaz NP)
Saffron Finch - @Mjly_jMZnjM		
Blue-black Grassquit - GnjMb bMjMjM M		(Porce area)
Ruddy-breasted - @ZXZMZx@unxualjMk b {M	« «	
Variable Seedeater - @unxualjMhxftb M		(Tatamá NP)
Gray Seedeater - @unxualjMb {Zk ZXbM		
Black-and-white - @ZXZMZx@unxualjM V nyM		(Porce area)
Yellow-bellied Seedeater - @unxualjM b xbhjly	« «	
Band-tailed Seedeater - t MjM ZI bMjMly		(Nevados NP)
Plain-colored Seedeater - t MjM ZI Mb nx MjM		(Nevados NP)
Paramo Seedeater - t MjM ZI Mb nk nVaxnM	« «	
Bananaquit - t nZxZUMjMZnjM		(Porce area)
Yellow-faced Grassquit - AbMjnjbMZY y	« «	
Buff-throated Saltator - @jMjMnxk Mlk y	" "	
Black-winged Saltator - @jMjMnxMbzI by		(La Romera)
Grayish Saltator - @jMjMnxVhZx jZYZI y		(Porce area)
Streaked Saltator - @jMjMnxy{xMbZV y		(Otún Q)
Masked Saltator - @jMjMnxVb V y	Near-threatened Vulnerable	(Río Blanco R)

SPARROWS & ALLIES: Emberezidae

Tanager Finch - 5xZn{axMlubyMxZk nl nuy	Near endemic Vulnerable	(Tatamá NP)
Yellow-throated Chlorospingus - t ajnxnyub` y_jjMb jMly		(Piha R)
Ashy-throated Chlorospingus - t ajnxnyub` yVMb jMly		(Río Blanco R)
Common Chlorospingus - t ajnxnyub` y_jMfnuZV y		(La Romera)
Dusky Chlorospingus - t ajnxnyub` yyZk b yM y		(Tatamá NP)
Black-headed Brushfinch - ° xxZk nl MbMjly	Near endemic	(Porce area)
Gray-browed Brushfinch - ° xxZk nl Mjly		(Río Blanco R)
Chestnut-capped Brushfinch - ° xxZk nl Ux I Zb VaM		§ §
Olive Finch - ° xxZk nl MjMjMjZbZuy	Near-threatened	(Tatamá NP)

Rufous-collared	- @lMn, Lnl n{xVatMMZI yly	(Río Blanco R)
White-naped Brushfinch	- ° {jMIZ{ZyMUb VaM	(La Romera)
Moustached Brushfinch	- ° {jMIZ{ZyMh_xZl M y' Near endemic	(Chicaque R)
Tricolored Brushfinch	- °° {jMIZ{Zy{xhjn	(Tatamá NP)
Slaty Brushfinch	- ° {jMIZ{ZyVaby{MIZ y	(Yellow-eared Parrot R)
Pale-naped Brushfinch	- ° {jMIZ{ZyUj b VaM	(Nevados NP)
Yellow-breasted Brushfinch	- ° {jMIZ{ZyjMb Va y	(Chicaque R)

CARDINALS & GROSBEAKS: Cardinalidae

Summer Tanager	- <bMI` Mx UxM	(Piha R)
Red-hooded Tanager	- <bMI` Mx UxZuy	(Río Blanco R)
White-winged Tanager	- <bMI` MZ Vhu{ZxM	(Piha R)
Crested Ant-Tanager	-) MUMx{MIM Endemic	(Tatamá NP)
Black-backed Grosbeak	- <aZ Vb yM xZnFZl {xy	(Chicaque R)
Rose-breasted Grosbeak	- <aZ Vb yj XnFbM y	(Porce area)

WOOD WARBLERS: Parulidae

Northern Waterthrush	- <Mi ZyM nfZUhxMIZl yly	(Montezuma farm)
Black-and-white Warbler	- 2 l bn{b{MFM	(Río Blanco R)
Tennessee Warbler	- 1Zbn{ajtulyuZx' x M	(Piha R)
Connecticut Warbler	- 5unxn byM ly	«- «-
Mourning Warbler	- (Zn{ajtulyuajMxZuatM	«- «-
American Redstart	- @Z{nuaMIM {bMjM	(Otún Q)
Tropical Parula	- @Z{nuaMIMUk k b	(Porce area)
Bay-breasted Warbler	- @Z{nuaMIMMx{MIZM	«- «-
Blackburnian Warbler	- @Z{nuaMIM yM	(Otún Q)
Yellow Warbler	- @Z{nuaMIMUz{ZVatM	(Piha R)
Black-crested Warbler	- 2 tbn{ajtuly b xnVxy{MIM	(Chicaque R)
Russet-crowned Warbler	- 2 tbn{ajtulyVhxn MIM	(La Romera R)
Three-striped Warbler	- E MjZ {Zx y{x xM y	" "
Canada Warbler	- f MxZj b M M M M x Zl yly	«- «-
Slate-throated Redstart	- 2 tbnUhx yk b bM y	(Río Blanco R)
Golden-fronted Redstart	- 2 tbnUhx ynx M y' Near endemic	(Chicaque R)

ICTERIDS: Icteridae

Russet-backed Oropendola	- <yMnVhjb yM y{b nl y	(Piha R)
Crested Oropendola	- <yMnVhjb yXZV k M y	(Porce area)
Mountain Cacique	- f M yVaxtynl n{ y	(Yellow-eared Parrot R)
Yellow-billed Cacique	- ° k UfVZxM yanjnyZxZ y	(Río Blanco R)
Yellow-tailed Oriole	- VZx yk Zynk ZjM	(Porce area)
Yellow-backed Oriole	- VZx yVaxtyMz	«- «-
Red-bellied Grackle	-) tunutxa yutxnatun` MxZx Endemic Vulnerable	(La Romera)

